## 論文

- Olmai T, Kuwahara K, Kabe I, et al.; Impact of a reduction in overtime hours on sleep duration among Japanese employees. Environ Occup Health Practice. 2020; 2: eohp.2020-0012-CM.
- https://www.jstage.jst.go.jp/article/eohp/advpub/0/advpub\_2020-0012-CM/\_article/-char/ja/
- ○Nanri A, Nakamoto K, Sakamoto N, et al.; Green tea consumption and influenza infection among Japanese employees. Eur J Clin Nutr. 2020 Nov 2. (共著)
- ○Fukunaga A, Hu H, Inoue Y, et al.; Diabetes, prediabetes, and suicide deaths in a Japanese working population. J Psychosom Res. 2020; 138:110246. (共著)
- ○Sakamoto N, Hu H, Nanri A, et al.; Associations of anemia and hemoglobin with hemoglobin A1c among non-diabetic workers in Japan. J Diabetes Investig. 2020; 11:719-725. (共著)
- ○Hu H, Tomita K, Kuwahara K, et al.; Obesity and risk of hearing loss: A prospective cohort study. Clinical Nutrition, 2020; 39; 870-5. (共著)
- Endo M, Inoue Y, Kuwahara K, et al.; BMI and Medically Certified Long-Term Sickness Absence Among Japanese Employees. Obesity, 2020; 28: 437-44. (共著)
- ○Xiang M, Hu H, <u>Imai T</u>, et al.; Association between anthropometric indices of obesity and risk of cardiovascular disease in Japanese men. J Occup Health, 2020; 62, e12098. (共著)
- OHasegawa M, Akter S, Hu H, et al.; Five-year cumulative incidence of overweight and obesity, and longitudinal change in body mass index in Japanese workers: The Japan Epidemiology Collaboration on Occupational Health Study. J Occup Health, 2020; 62, e12095. (共著)
- OKuwahara K, Endo M, Nishiura C, et al.; Smoking cessation after long-term sick leave due to cancer in comparison with cardiovascular disease: Japan Epidemiology Collaboration on Occupational Health Study. Ind Health. 2020;58:246-253. (共著)
- Olnoue Y, Nomura S, Nishiura C, et al.; Loss of working life years due to mortality, sickness absence, or illhealth retirement: A comprehensive approach to estimating disease burden in the workplace. J Epidemiol. 2020; JE20190332. (共著)